Mini-course on Sparse estimation off-the-grid Introduction

Clarice Poon

• Session 1: Introduction to sparse estimation as optimisation over the space of measures

• Session 2: Algorithms

• Session 3: Super resolution and compressed sensing guarantees

Outline

Sparse Estimation **Recovering point wise sources from low resolution data**

Let $\mathscr{X} \subseteq \mathbb{R}^d$ and let $\phi : \mathscr{X} \to \mathscr{H}$ where \mathscr{H} is a Hilbert space.

Recover $a_i \in \mathbb{R}$ and $x_i \in \mathcal{X}$ given y =



$$\sum_{j=1}^{s} a_{j} \phi(x_{j})$$



The space of Radon measures $\mathscr{M}(\mathscr{X})$ is the dual of

$$C_0(\mathcal{X}) = \left\{ f \in C(\mathcal{X}) : f \right\}$$

View $\mu \in \mathcal{M}(\mathcal{X})$ as linear functional on $C_0(\mathcal{X})$:

•For
$$f \in L^1(\mathcal{X})$$
, define μ by $\langle \phi, \mu \rangle = \int \phi(x) f(x) dx$
•For $\mu = \sum_j a_j \delta_{x_j}$, $\langle \phi, \mu \rangle = \sum_j \phi(x_j) a_j$

Radon measures







Linear inverse problem

Consider a measure μ on $\mathcal{X} \subseteq \mathbb{R}^d$





Observe linear measurements:

Define:
$$\Phi \mu = \int_{\mathcal{X}} \phi(x) d\mu(x)$$

ſ

 $\phi(x) \in \mathcal{H}$ where $\phi : \mathcal{X} \to \mathcal{H}$

Observe: $y = \Phi \mu + noise$

NB:
$$\Phi \mu_{a,x} = \sum_{i=1}^{n} a_i \phi(x_i)$$



Signal/image processing

Deconvolution:

$$\phi(x) = \tilde{\phi}(\cdot - x) \in L^2(\mathbb{R}^d)$$

e.g. $\tilde{\phi}(x) = \exp(|x - \cdot|^2/\sigma)$

Laplace:

 $\phi(x) = \exp(-\langle x, \cdot \rangle) \in L^2(\mathbb{R}^d_+)$



Fourier:

 $\phi(x) = (\exp(kx\sqrt{-1}))_{k=-f_c,...,f_c} \in \mathbb{C}^{2f_c+1}$











 θ_{γ}

$\theta = T_1/T_2$ representing tissue type $\phi(\theta) =$ Block response of each tissue

Time series data $Y = (y^{\nu})$



Quantitative MRI



There can be more than 1 tissue type in each image voxel (so n > 1).





Mixture models



Density estimation with sketching





Goal: recover *a*, *x* from

Given samples t_1, t_2, \ldots, t_n iid from from density: $\bar{\xi}(t) = \sum_{j=1}^{s} a_{j}\xi(x_{j}, t) = \int \xi(x, t) d\mu_{a,x}(x)$ i=1

[Gribonval et al 2017]

Sketch using functions g_{ω_k} : $y_k = \frac{1}{n} \sum_{j=1}^n g_{\omega_k}(t_j), k \in [m]$

$$y_k \approx \int g_{\omega_k}(t) \bar{\xi}(t) dt = \int_{\mathcal{X}} \underbrace{\int g_{\omega_k}(t) \xi(x, t) dt}_{\phi_{\omega_k}(x)} d\mu_{a, x}(x)$$



Multi-laye



Non-convex

$$\min_{a,z,b} \sum_{i} |f_{a,z,b}(t_i) - y_i|^2$$

Convex

 $\min_{\mu \in \mathcal{M}(\mathcal{X})} \| y - \Phi \mu \|^2$

$$f_{a,z,b}(t) = \sum_{i=1}^{n} a_i \rho(\langle z_i, t \rangle + b_i) \in \mathbb{R}^n$$

$$[f_{a,z,b}(t_i)]_i = \Phi \mu = \int_{\mathbb{R}^d} \phi(x) d\mu(x)$$
$$\phi(x) = \left[\rho(\langle z, t_i \rangle + b) \right]_{i=1,\dots,N} \quad \mu = \sum_{i=1}^n a_i \, \delta_{(z_i)}$$

Linear operator







$\mathcal{M}(\mathcal{X})$ is a Banach space with norm $\|\mu\|_{TV}$ $\|\mu\|_{TV} = \sup\left\{ \int f(x) d\mu(x) : f \in C_0(\mathcal{X}), \|\|f\|_{\infty} \le 1 \right\}$

 $f \in L^1(\mathcal{X}), d\mu(x) = f(x)dx \longrightarrow \|\mu\|_{TV} = \|f(x)\|dx$ $\mu = \sum_{j} a_{j} \delta_{x_{j}} \longrightarrow \|\mu\|_{TV} = \sum_{j} |a_{j}|$

The extremal points of $\{\mu : \|\mu\|_{TV} \le 1\}$ are $\{\delta_x : x \in \mathcal{X}\}$

Total variation



 $P_{\lambda}(y)$

Fisher-Jerome (1973): If $\phi(x) \in \mathbb{R}^m$ with ϕ continuous, then there exists a solution to $P_{\lambda}(y)$ with at most *m* Diracs.

 $\inf_{\mu \in \mathscr{M}(\mathscr{X})} \|\mu\|_{TV} \quad \text{s.t.} \quad \Phi \mu = y \,.$ $P_0(y)$

The Beurling-Lasso $\inf_{\mu \in \mathcal{M}(\mathcal{X})} \lambda \|\mu\|_{TV} + \frac{1}{2} \|\Phi\mu - y\|^2$

Relaxation for any K: $\inf_{a,x} \lambda \sum_{j=1}^{K} |a_j| + \frac{1}{2} \|\sum_{j=1}^{K} \phi(x_j)a_j - y\|^2 \ge \inf P_{\lambda}(y)$

The relaxation is tight when $K \ge m$

[Beurling (1973)]

[De Castro and Fabrice (2012)]

[Candès and Fernandez-Granda (2012)]

[Duval and Peyré (2015).]



The Beu





- Optimisation is over the space of measures (not just Diracs) with no a-priori choice on the number of spikes.
- This is a convex problem, with strong recovery guarantees.
- Some non-convex problems can be placed into this framework

$$\frac{1}{2} \|\Phi\mu - y\|^2$$

The Lasso: Given y = Xa, $y \in \mathbb{R}^m$, $X \in \mathbb{R}^{m \times n}$, to recover a sparse vector $a \in \mathbb{R}^n$ $\min_{a \in \mathbb{R}^n} \frac{1}{2\lambda} \|Xa - y\|^2 + \|a\|_1$

Questions

- When is $\mu_0 = \sum_j a_j \delta_{x_j}$ an exact solution to $(P_0(y))$?
- Are solutions to $P_{\lambda}(y)$ stable to noise?
- Numerical algorithms in the infinite dimensional space?
- Inder what conditions do we recover the exact number of spikes?
- ${\ensuremath{\, \bullet }}$ Compressed sensing if Φ is a random operator, how many measurements to recover?

Optimality conditions $\mu_* \in \operatorname{argmin}_{\mu} F(\mu) \quad \clubsuit \quad \nabla F(\mu_*) = 0$

But $\|\mu\|_{TV}$ is not differentiable. Need to consider its **sub-differential**.

Let $\Psi: U \to \mathbb{R}$ be a convex function, its **sub-differential** is: $\partial \Psi(\mu) = \left\{ p \in U^* : \forall \hat{\mu}, \Psi(\hat{\mu}) \ge \Psi(\mu) \right\}$

$$\mu) + \langle p, \hat{\mu} - \mu \rangle \bigg\}$$





Optimality conditions

Equivalent characterization for $\|\mu\|_{TV}$: $\partial \|\mu\|_{TV} = \{f \in C(\mathcal{X}) : \|f\|_{\infty} \le 1, \langle f, \mu \rangle = \|\mu\|_{TV} \}$



 $\partial \|\mu_{a,x}\|_{TV} = \left\{ f \in C(\mathcal{X}) : \left\{ \begin{array}{l} \|f\|_{\infty} \leq 1 \\ \forall i, \ f(x_i) = \operatorname{sign}(a_i) \end{array} \right\} \right\}$ $\mu_{a,x} = \sum_{i}^{n} a_i \delta_{x_i}$





Optimality conditions

For convex problem $\min_{x} F(x)$, minimiser iff $0 \in \partial F(x)$

$$\mu_{\lambda} \in \underset{\mu \in \mathcal{M}(\mathcal{X})}{\operatorname{argmin}} \lambda \|\mu\|_{TV} + \frac{1}{2} \|\Phi\mu - y\|^{2}$$



•
$$0 \in \partial \|\mu_{\lambda}\|_{TV} + \frac{1}{\lambda} \Phi^*(\Phi\mu_{\lambda} - y)$$

$$\eta_{\lambda} := -\frac{1}{\lambda} \Phi^*(\Phi \mu_{\lambda} - y) \in \partial \|\mu_{\lambda}\|_{TV}$$

$\operatorname{Supp}(\mu_{\lambda}) \subset \{x : |\eta_{\lambda}(x)| = 1\}$

The *dual certificate* η_{λ} certifies the support of μ_{λ}



Convex duality

 $\min_{\mu \in \mathcal{M}(\mathcal{X})} \|\mu\|_{TV} + \frac{1}{2\lambda} \|\Phi\mu - y\|^2$ Primal:

Dual:
$$\sup_{\|\Phi^*p\|_{\infty} \le 1} \langle p, y \rangle - \frac{\lambda}{2} \|p\|^2 \qquad (D)$$



Convex duality

Dual:
$$\sup_{\|\Phi^*p\|_{\infty} \le 1} \langle p, y \rangle - \frac{\lambda}{2} \|p\|^2 = -\frac{\lambda}{2} \|p - y/\lambda\|^2 + \frac{1}{\lambda} \|y\|^2$$

- $D_{\lambda}(y)$ is the projection onto a convex set. So, it has a unique solution.
- There is strong duality. $\inf P_{\lambda}(y) = \sup D_{\lambda}(y)$
- When $\lambda > 0$, solutions to $P_{\lambda}(y)$ and $D_{\lambda}(y)$ exist.

The noiseless problem

Primal: min $\|\mu\|_{TV}$ s.t. $\Phi\mu = y$ $\mu \in \mathcal{M}(\mathcal{X})$

• When $\lambda = 0$, only existence of solutions to $P_0(y)$ is guaranteed (unless \mathcal{H} is finite).

Projection onto convex set

• If $\mathcal{H} = \mathbb{R}^n$, optimise over finite vector space but with infinite constraints.





Convex duality



If $p_{\lambda} = \operatorname{argmax} D_{\lambda}(y)$ and $\eta_{\lambda} = \Phi^* p_{\lambda}$, then $\eta_{\lambda} \in \partial \|\mu_{\lambda}\|_{TV}$ means that $\operatorname{Supp}(\mu_{\lambda}) \subset \{x : |\eta_{\lambda}(x)| = 1\}$

Solutions to $D_0(\Phi\mu_0)$ can tell us about the structure of $\mu_{\lambda} \in \min P_{\lambda}(\Phi\mu_0 + w)$

$$\mu_0 \text{ solves } P_0(y) \text{ and } p_0 \text{ solves } D_0(y)$$



Theorem: If $\mu_{a,x} = \sum a_j \delta_{x_i}$ and $y = \Phi \mu_{a,x}$ and there exists *p* such that • $\eta := \Phi^* p$ satisfies $|\eta(x)| < 1$ for all $x \notin \{x_i\}$ • $\eta(x_i) = \operatorname{sign}(a_i)$ for all *i*. $(\phi(x_i))_i$ are linearly independent. \bigcirc

Then, $\mu_{a,x}$ is the unique solution to $P_0(y)$

So, any two solutions take the form: μ

We must have $a_i = \hat{a}_i$ since $\Phi \hat{\mu} = \Phi \mu$ and $\phi(x_i)$ are linearly independent.

Uniqueness



Proof: by the primal-dual relationships, any solution has support contained in $\{x_i\}_i$

$$= \sum_{i} a_{i} \delta_{x_{i}} \text{ and } \hat{\mu} = \sum_{i} \hat{a}_{i} \delta_{x_{i}}$$

Stability

Theorem [Azais De Castro & Gamboa (2015)] Suppose we observe $y = \Phi \mu_{a,x} + w$ with $||w|| \le \epsilon$. In addition to conditions of previous theorem, suppose $\eta = \Phi^* p$ satisfies i) ii) Then, choosing $\lambda \sim \epsilon/\|p\|$, any solution $\hat{\mu}$ to $P_{\lambda}(y)$ satisfies $c_0 |\hat{\mu}| (\mathcal{X} \setminus \bigcup_i B(x_i, r)) + c_2 \sum_i \int_{B(x_i, r)} ||x - x_i||^2 \mathrm{d} |\hat{\mu}| (x) \lesssim \epsilon ||p||$ amplitudes outside neighbourhood of true support is small

- $|\eta(x)| \le 1 c_2 ||x x_i||^2$ for all $x \in B(x_i, r)$ $|\eta(x)| < 1 - c_o$ for all $x \notin \bigcup_i B(x_i, r)$

Cluster around true support



Stability

Theorem [Azais De Castro & Gamboa (2015)] Suppose we observe $y = \Phi \mu_{a,x} + w$ with $||w|| \le \epsilon$. In addition to conditions of previous theorem, suppose $\eta = \Phi^* p$ satisfies i) ii) Then, choosing $\lambda \sim \epsilon/\|p\|$, any solution $\hat{\mu}$ to $P_{\lambda}(y)$ satisfies $c_0 |\hat{\mu}| (\mathscr{X} \setminus \bigcup_i B(x_i, r)) + c_2 \sum_i \int_{B(x_i, r)} ||x - x_i||^2 \mathrm{d} |\hat{\mu}| (x) \lesssim \epsilon ||p||$

If $\eta \in \text{Im}(\Phi^*)$ satisfies (i) and (ii), then we say that it is nondegenerate.

- $|\eta(x)| \le 1 c_2 ||x x_i||^2$ for all $x \in B(x_i, r)$ $|\eta(x)| < 1 - c_0$ for all $x \notin \bigcup_i B(x_i, r)$
- $W_2^2(\sum_{i} \hat{A}_j \delta_{x_j}, |\hat{\mu}|) \lesssim \epsilon \|p\| \quad \text{and} \max_j |a_j \hat{a}_j| \lesssim \epsilon \|p\|$ $\hat{A}_i = |\hat{\mu}| (B(x_i, r)) \quad \hat{a}_j = \hat{\mu}(B(x_j, r))$



Candidate for a dual certificate

Define:

$$K(x_1, x_2) = \langle \phi(x_1), \phi(x_2) \rangle$$

Want: $\eta(x_i) = \operatorname{sign}(a_i)$ and $\eta'(x_i) = 0$

Computed η and check if $|\eta(x)| < 1$ for all $x \notin \{x_i\}$.





$$\eta_C(x) = \sum_{i=1}^n u_i K(x_i, x) + \sum_{i=1}^n v_i \partial_1 K(x_i, x)$$







Recovery under minimal separation

Candès and Fernandez-Granda (2012): Let $\phi(x) = (\exp(2\pi\sqrt{-1}kx)_{|k| \le f_c})$ if $\min_{i \neq j} |x_i - x_j| \ge \frac{C}{f_c}$, then η_C is non-degenerate. So, we have stable recovery.

Necessary: If
$$|x_1 - x_2| < \frac{1}{f_c}$$
 then μ



Recovery under minimal separation



translation-invariant

What kind of minimum separation condition to impose for non-translation invariant kernel?



Not translation-invariant



Fisher-Rao distance

Fisher metric: $g_x := \partial_1 \partial_2 K(x, x') =$

Fisher-Rao geodesic distance: $d_g(x,$



$$= [\nabla \phi(x)] [\nabla \phi(x')]^\top \in \mathbb{R}^d$$

$$(x') := \inf_{\gamma:x \to x'} \int_0^1 \sqrt{\langle g_{\gamma(t)} \gamma'(t), \gamma'(t) \rangle} dt$$

Interpretation:

 $x \mapsto \phi(x)$ embeds \mathscr{X} into the sphere in \mathscr{H} and

$$d_g(x, x') = \inf_{\substack{\gamma:\phi(x)\to\phi(x')}} \int_0^1 \|\gamma'(t)\|_{\mathcal{H}} dt$$

Examples

Poon, Keriven and Peyre (2019): If $\min_{i \neq j} d_g(x_i, x_j) \ge \Delta_{s,K}$, then η_C is nondegenerate.

Gaussian	Fourier	Laplace
$\phi(x) \propto \exp(-\ x - \cdot\ _{\Sigma}^2)$	$\phi(x) = (\exp(2\pi\sqrt{-1}kx))_{\ k\ _{\infty} \le f_c}$	$\phi(x) \propto \exp(-x \cdot)$
$g_x = \Sigma$	$g_x = f_c I$	$g_x = \operatorname{diag}(1/x_i)$
$d_g(x, x') = \ x - x'\ _{\Sigma}$	$d_g(x, x') \propto f_c x - x' _2$	$d_g(x, x') = \sqrt{\sum_{i} \log(x_i) - \log(x'_i) ^2}$
$\Delta = \sqrt{\log(s)}$	$\Delta = \sqrt{d\sqrt{s}}$	$\Delta = d + \log(ds)$

_ ,	

Exact recovery in the noiseless setting Provided that spikes are sufficiently separated: - Stable recovery in the noisy setting.

Summary

on
$$\lim_{a,x} \lambda \sum_{j=1}^{K} |a_j| + \frac{1}{2} \|\sum_{j=1}^{K} \phi(x_j) a_j - \sum_{j=1}^{K} \phi(x_j) a_j - \sum$$

To assess the recovery of $m_{a,x}$, Find $\eta = \Phi^* p \in C(\mathcal{X})$ such that $\eta(x_i) = \operatorname{sign}(a_i)$ and $|\eta(x)| < 1$ for all $x \notin \{x_i\}$

References

General theory for the Blasso:

- •Candès, E. J., & Fernandez-Granda, C. (2014). Towards a mathematical theory of super-resolution. Communications on pure and applied Mathematics, 67(6), 906-956.
- •Azais, J. M., De Castro, Y., & Gamboa, F. (2015). Spike detection from inaccurate samplings. Applied and Computational Harmonic Analysis, 38(2), 177-195.
- •Bredies, K., & Pikkarainen, H. K. (2013). Inverse problems in spaces of measures. ESAIM: Control, Optimisation and Calculus of Variations, 19(1), 190-218.
- Duval, V., & Peyré, G. (2015). Exact support recovery for sparse spikes deconvolution. Foundations of Computational *Mathematics*, *15*(5), 1315-1355.
- Poon, C., Keriven, N., & Peyré, G. (2021). The geometry of off-the-grid compressed sensing. Foundations of Computational Mathematics, 1-87

A few references for applications

- and its application to super-resolution microscopy. *Inverse Problems*, 36(1), 014001. learning with random feature moments. *Mathematical Statistics and Learning*, 3(2), 113-164 Journal of Machine Learning Research, 18(1), 629-681.
- Denoyelle, Q., Duval, V., Peyré, G., & Soubies, E. (2019). The sliding Frank–Wolfe algorithm • Gribonval, R., Blanchard, G., Keriven, N., & Traonmilin, Y. (2021). Compressive statistical •Bach, F. (2017). Breaking the curse of dimensionality with convex neural networks. The
- Golbabaee & Poon (2022). An off-the-grid approach to magnetic resonance fingerprinting. Inverse Problems (to appear).

